

Scattering of 8 He on 208 Pb at 22 MeV

G. Marquínez-Durán, A. M. Sánchez-Benítez, I. Martel, L. Acosta, K. Rusek, M. A. G. Álvarez, R. Berjillos, M. J. G. Borge, A. Chbihi, C. Cruz, M. Cubero, J. A. Dueñas, J. P. Fernández-García, B. Fernández-Martínez, J. L. Flores, J. Gómez-Camacho, N. Keeley, J. A. Labrador, M. Marqués, A. M. Moro, M. Mazzocco, A. Pakou, V. V. Parkar, N. Patronis, V. Pesudo, D. Pierroutsakou, R. Raabe, R. Silvestri, N. Soic, . Standylo, I. Strojek, O. Tengblad, R. Wolski, and A. H. Ziad

Citation: *AIP Conference Proceedings* **1541**, 175 (2013); doi: 10.1063/1.4810834

View online: <http://dx.doi.org/10.1063/1.4810834>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/1541?ver=pdfcov>

Published by the [AIP Publishing](#)

Scattering of ^8He on ^{208}Pb at 22 MeV

G. Marquín-Durán^{*}, A.M. Sánchez-Benítez^{*}, I. Martel^{*}, L. Acosta^{*,†},
K. Rusek^{**}, M.A.G. Álvarez^{‡,§}, R. Berjillos^{*}, M.J.G. Borge[¶], A. Chbihi^{||},
C. Cruz[¶], M. Cubero[¶], J.A. Dueñas^{*}, J.P. Fernández-García[§],
B. Fernández-Martínez[‡], J.L. Flores^{††}, J. Gómez-Camacho^{‡,§}, N. Keeley^{‡‡},
J.A. Labrador[‡], M. Marqués^{§§}, A.M. Moro[§], M. Mazzocco^{¶¶}, A. Pakou^{***},
V.V. Parkar^{*}, N. Patronis^{***}, V. Pseudo[¶], D. Pierroutsakou^{†††}, R. Raabe^{‡‡‡},
R. Silvestri^{††††}, N. Soic^{§§§}, Ł. Standylo^{‡‡}, I. Strojek^{‡‡}, O. Tengblad[¶],
R. Wolski^{¶¶¶} and A.H. Ziad[‡]

^{*}Depto. de Física Aplicada, Universidad de Huelva, 21071 Huelva, Spain

[†]Laboratori Nazionali del Sud, INFN, Via Santa Sofia 62, 95123, Catania, Italy

^{**}Heavy Ion Laboratory, University of Warsaw

[‡]Centro Nacional de Aceleradores, 41092, Sevilla, Spain

[§]Depto. de Física Atómica, Molecular y Nuclear, Universidad de Sevilla, 41080 Sevilla, Spain

[¶]Instituto de Estructura de la Materia, CSIC, 28006 Madrid, Spain

^{||}GANIL, CEA and IN2P3-CNRS, B.P. 5027, 14076 Caen cedex, France

^{††}Depto. de Ingeniería Eléctrica, Universidad de Huelva, 21071 Huelva, Spain

^{‡‡}National Centre for Nuclear Research, A.Soltana 7, 05-400 Otwock, Poland

^{§§}Laboratoire de Physique Corpusculaire, 14050 Caen cedex, France

^{¶¶}Dip. di Fisica and INFN, Università di Padova, via F. Marzolo 8, 35131 Padova, Italy

^{***}Department of Physics, University of Ioannina, 45110 Ioannina, Greece

^{†††}INFN - Sezione di Napoli, via Cintia, 80126 Napoli, Italy

^{‡‡‡}Instituut voor Kern-en Stralingsfysica, K.U. Leuven, Belgium

^{§§§}Rudjer Boskovic Institute, Bijenicka 54, 10000 Zagreb, Croatia

^{¶¶¶}Flerov Laboratory of Nuclear Reactions, JINR, Dubna 141980, Russian Federation

Abstract. The skin nucleus ^8He is investigated by measuring the angular distribution of the elastically scattered ^8He and the $^6,4\text{He}$ fragments produced in the collision with a ^{208}Pb target at 22 MeV, just above the Coulomb barrier. The experiment was carried out at SPIRAL/GANIL in 2010. Here we present preliminary results for the elastic scattering.

Keywords: Direct reactions, Elastic scattering, ^8He

PACS: 01.30.Cc, 13.85.Dz, 24.10.Ht, 25.10.+s

INTRODUCTION

The near-barrier scattering of ^8He with heavy targets remains still not very well known. Only a few scattering data sets are available nowadays [1, 2]. The skin nucleus ^8He presents the largest N-Z asymmetry of the particle-stable nucleus and its particular features makes it an attractive field of study. This work is focused in the experimental results for the elastic scattering up to 95° , for the $^8\text{He}+^{208}\text{Pb}$ system at 22 MeV.

EXPERIMENTAL SETUP

The experiment was performed at SPIRAL/GANIL. The experimental setup consisted of a portable reaction chamber where the array of detectors was mounted, a set of collimators and a beam diagnostics system. The detection system, GLORIA (GLOBal Reaction Array), consists of 12 DSSSD detectors arranged in 6 particle telescopes (40 μ m, 1mm) covering a continuous angular range between 15° and 165°.

EXPERIMENTAL RESULTS FOR THE ELASTIC SCATTERING

The experimental results obtained are presented in Fig. 1. The three first telescopes, covering up to 95°, are plotted separately in order to show the existing matching between them. The elastic cross section decreases in a smooth way losing the Coulomb-nuclear rainbow, characteristic for light stable nuclei.




FIGURE 1. Angular distribution of the elastic cross section for the system ${}^8\text{He}+{}^{208}\text{Pb}$ at 22 MeV. The two forward and the bottom telescopes are represented separately.

The experimental data is compared with an OM calculation in which the optical potential derived from the elastic scattering of ${}^8\text{Li}+{}^{208}\text{Pb}$ [3] is used.

CONCLUSIONS

The elastic scattering of ${}^8\text{He}+{}^{208}\text{Pb}$ at 22 MeV has been measured at SPIRAL-GANIL using GLORIA, a new detector developed at the University of Huelva. The angular distribution of the elastic channel decreases in a smooth way losing the Coulomb-nuclear rainbow.

REFERENCES

1. A. Lemasson *et al.*, *Phys. Rev.* **C82**, 044617 (2010).
2. A. Lemasson *et al.*, *Phys. Lett.* **B697**, 454 (2011).
3. J.J. Kolata *et al.*, *Phys. Rev.* **C65**, 054616 (2002).